

model. Analyses limiting the dementia outcome to AD alone (45 cases) had essentially the same results. **Conclusions:** These findings suggest that WMH may interfere with cognitive function but may not be sufficient to produce dementia. In persons whose cognitive abilities are already impaired, a global measure of brain atrophy is more predictive of conversion to dementia. Persons at increased risk of stroke are also at increased risk of dementia.

P2-358 COGNITIVE PROCESSING SPEED AND MYELIN BREAKDOWN IN OLDER INDIVIDUALS

George Bartzokis, Po H. Lu, Kathleen Tingus, Nancy Edwards, Christopher A. Lieu, Mario F. Mendez, Jim Mintz, Jeffrey L. Cummings, *UCLA, Los Angeles, CA, USA. Contact e-mail: gbar@ucla.edu*

Background: The process of myelination peaks in middle age followed by myelin breakdown and loss. Investigations of information processing speed performance across the lifespan reveal a trajectory that parallels myelination and subsequent myelin breakdown. Myelin breakdown may result in a progressive “disconnection” of widely distributed neural networks and may underlie age-related cognitive decline and Alzheimer’s disease (AD). **Objective(s):** This study tests the “myelin model”/hypothesis that in older individuals, myelin breakdown in late-myelinating regions is associated with slowed processing speed and contributes to the continuum of cognitive decline that ultimately results in AD. **Methods:** Healthy older (>55 years) individuals (N=92) and 8 subjects with Alzheimer’s disease (AD) had neurocognitive testing. The late-myelinating frontal lobe white matter (Fwm) as well as early- and later-myelinating regions of the corpus callosum, the splenium (Swm) and genu (Gwm) respectively, were assessed using MRI and transverse relaxation rates (R_2) were calculated. R_2 is an indirect measure of white matter structural integrity; it declines with age-related myelin breakdown and is significantly lower in AD. **Results:** As hypothesized, cognitive processing speed tasks (Trails A and Digit Symbol) were significantly associated with R_2 in late-myelinating regions (Fwm, $p<.0001$ and Gwm, $p<.004$) but not in the early myelinating Swm region ($p>.2$). **Conclusions:** These data suggest that myelin breakdown in healthy older individuals underlies the age-related cognitive decline that ultimately results in AD. MRI measures of myelin breakdown and cognitive measures may be useful in AD primary prevention studies.

P2-359 PREDICTING COGNITIVE DECLINE IN MCI AND HEALTHY AGING WITH STRUCTURAL AND PERFUSION MRI AND MRS

Linda L. Chao¹, Norbert Schuff¹, Audrey Duarte², Shannon T. Buckley¹, John Kornak¹, Joel H. Kramer¹, Kristine Yaffe¹, Bruce L. Miller¹, Michael W. Weiner¹, ¹University of California, San Francisco, San Francisco, CA, USA; ²Medical Research Council, Cambridge, United Kingdom. *Contact e-mail: llchao@radmail.ucsf.edu*

Background: Potentially effective treatments for Alzheimer’s disease (AD) are beginning to enter human trials. Thus, it is important to develop measurements that can predict cognitive decline in mildly impaired subjects so that individuals at greatest risk for AD can be targeted when preventative treatments do become available. **Objective:** To determine the regional pattern of structural, perfusion, and metabolic changes in the brain that predict cognitive decline. **Methods:** Fifty-five subjects (75.1 ± 7 years old) were assessed with neuropsychological tests at baseline and 1.2 ± 0.2 years later. Forty-eight subjects were MCI patients and 7 were healthy controls. Multiple regression was used to investigate the relationship between baseline cognition, volumetric structural MRI data (available for 45 subjects), and ¹H MRS data (42 subjects) and cognitive decline. Separate regressions were computed for change in MMSE, CDR, CDR sum of boxes, and different California Verbal Learning Test (CVLT) measures. Baseline measures were dependent variables, cognitive change scores were independent variables, and age, group, and test interval were nuisance variables in the analyses. The relationship between baseline cerebral per-

fusion, as determined by arterial spin labeled MRI (33 subjects) and cognitive decline were assessed with SPM2. **Results:** Baseline CVLT measures were negatively associated with in MMSE, CDR, and CDR sum of boxes change scores (i.e., lower baseline scores associated with greater decline, $p<0.05$). Baseline frontal and parietal gray matter (GM) volumes were also negatively associated with MMSE decline ($p<0.01$). Adding baseline frontal and parietal GM volumes to baseline CVLT measures significantly improved (p 0.03) the linear model. Baseline posterior cingulate N-acetylaspartate (NAA)/creatinine (Cr) ratio was negatively associated with decline in CDR sum of boxes ($p<0.05$). However, adding NAA/Cr ratio to baseline CVLT measures did not improve the linear model for predicting CDR sum of boxes decline. Voxel-by-voxel analyses revealed a negative relationship between baseline cerebral perfusion in the posterior cingulate and subsequent decline on CVLT short delay cued recall, even after accounting for baseline CVLT performance (see Figure 1). **Conclusions:** These preliminary results suggest that arterial spin labeled MRI and volumetric structural MRI may add predictive value to neuropsychological testing for predicting cognitive decline.

Figure 1. Regions showing significant correlations between perfusion and subsequent decline on CVLT short-delay cued recall, $p<0.05$, corrected for multiple comparisons.




P2-360 CORTICAL THINNING IN ALZHEIMER’S DISEASE AND FRONTOTEMPORAL DEMENTIA

Norbert Schuff¹, Antao Du¹, Howard Rosen², Maria Luisa Gorno-Tempini², Joel Kramer², Bruce L. Miller², Michael W. Weiner¹, ¹VAMC/UCSF, San Francisco, CA, USA; ²UCSF, San Francisco, CA, USA. *Contact e-mail: norbert.schuff@radiology.ucsf.edu*

Background: Volumetric MRI studies show characteristic patterns of brain atrophy in Alzheimer’s disease (AD) and frontotemporal dementia (FTD); however, an overlap between the groups remains. In contrast to general brain tissue loss, pathological studies reported predominantly loss of cortical layers in dementia, resulting in cortical thinning. Recently, MRI studies have used cortical thickness to better characterize brain alterations in dementia. **Objectives:** 1) To compare the regional patterns of cortical thinning in AD and FTD; 2) To determine if cortical thickness is a better measure than cortical volume to differentiate between normal aging, AD and FTD. **Methods:** 23 cognitive normal (CN) subjects, 21 AD and 19 FTD patients had MRI at 1.5T. Measurements of cortical thickness and volume were performed using FreeSurfer (<http://surfer.nmr.mgh.harvard.edu>). Regional differences in cortical thickness between the groups were tested vertex-by-vertex. The powers of cortical thickness and volume measures in frontal, parietal and temporal lobes to classify the groups were compared using logistic regression and receiver operator characteristics (ROC) analyses. **Results:** The figure depicts statistical maps of differences in cortical thinning between the groups. Compared to CN, AD was associated with thinner cortices in temporal, parietal and frontal regions, pre-cuneus and posterior cingulate ($p<0.001$), while FTD was associated with thinner cortices in primarily frontal and temporal regions ($p<0.001$). Com-

pared to FTD, AD was associated with thinner cortices in parietal regions, precuneus and posterior cingulate ($p < 0.01$). Compared to AD, FTD was associated with thinner cortices in orbitofrontal regions ($p < 0.01$). In general, cortical thickness provided no significant advantage over the corresponding volumes in classifying AD from CN and FTD. However, cortical thickness of parietal and temporal lobes (area under the ROC curve (AUC): 0.70-0.95 95% confident interval) showed a trend ($p = 0.06$) to better separate FTD from CN than the corresponding volume (AUC: 0.54-0.86). **Conclusions:** The characteristic patterns of cortical thinning in AD and FTD seen on MRI are consistent with pathological findings. Furthermore, cortical thinning may be a more prominent feature than volume loss for separating FTD from normal aging.



P2-361


DIFFERENT PATTERNS OF GRAY AND WHITE MATTER ATROPHY IN ALZHEIMER'S DISEASE AND SUBTYPES OF FRONTOTEMPORAL LOBAR DEMENTIA

Linda L. Chao, Norbert Schuff, Erin Clevenger, Maria L. Gorno-Tempini, Howard J. Rosen, Joel H. Kramer, Kristine Yaffe, Bruce L. Miller, Michael W. Weiner, *University of California, San Francisco, San Francisco, CA, USA. Contact e-mail: llchao@radmail.ucsf.edu*

Background: Alzheimer's disease (AD) and frontotemporal lobar degeneration (FTLD) are neurodegenerative diseases characterized by the progressive loss of cerebral tissue. Although structural magnetic resonance imaging (MRI) has been used extensively to investigate AD, and to a lesser extent FTLD, few studies have distinguished between the patterns of gray matter (GM) and white matter (WM) atrophy in these diseases. **Goals:** To identify the pattern of WM atrophy in two subtypes of FTLD—frontotemporal dementia (FTD) and semantic dementia (SD) and in AD relative to normal controls (CN). To compare the patterns of WM and GM atrophy in FTD, SD, and AD. **Methods:** Twelve FTD, 13 SD, 15 AD patients, and 24 cognitively normal subjects (CN) were studied with volumetric MRI. Regional GM and WM in each group were classified automatically from high resolution T1- and T2-weighted MRI using Expectation-Maximization Segmentation (EMS). Data were analyzed with analyses of covariance, with age and the white matter lesion volumes as covariates. **Results:** Figure 1 shows the Z-scores of GM and WM volumes by lobe for each patient group. Relative to CN, AD patients had significant GM atrophy in the parietal and temporal lobes ($p < 0.001$). Although there was a trend for

WM atrophy in the parietal lobe, there were no significant overall WM volume differences between AD and CN. FTD patients had significant GM and WM atrophy in the frontal lobe relative to CN ($p < 0.001$). There were trends of GM atrophy in the parietal and temporal lobes, but no WM atrophy in these regions. Compared to CN, SD patients had significant GM atrophy in both temporal and frontal lobes ($p < 0.001$), but significant WM atrophy only in the temporal lobe ($p < 0.0001$). **Conclusions:** These results suggest that different types of dementias are associated with different patterns of WM atrophy and that the pattern of WM atrophy differs by lobe. Whether WM atrophy in these diseases indicates general neuritic dystrophy, involving both neuron cell bodies and axons, or Wallerian degeneration remains to be explored.

Figure 1. Z scores of GM and WM volume in the different lobes for (A) AD patients, (B) FTD patients, and (C) SD patients.



P2-362

HIPPOCAMPAL ACTIVATION IN MCI PREDICTS SUBSEQUENT COGNITIVE DECLINE

Saul L. Miller¹, Julie Bates¹, Deborah Blacker¹, Reisa Sperling^{1,2}, Bradford C. Dickerson^{1,2}, *¹Massachusetts General Hospital, Charlestown, MA, USA; ²Brigham and Women's Hospital, Boston, MA, USA. Contact e-mail: smiller4@partners.org*

Background: We previously reported that fMRI hippocampal activation is greater in subjects with mild cognitive impairment (MCI) who subsequently demonstrate further cognitive decline. **Objective:** We investigated whether fMRI hippocampal activation predicts the degree of subsequent cognitive decline in MCI, and whether or not it predicts this degree of decline even after controlling for clinical predictors. **Methods:** Subjects included 27 older participants (ages 65-86) in a longitudinal memory and aging study for whom an average of 4.5 years (S.D. 1.1) of follow-up